## **Amendments to the Claims**

The following listing of claims will replace all prior versions, and listings, of claims in the application:

## Listing of Claims:

- 1. (Currently amended) A method of treating primary cancer which comprises administering to a patient in need of such treatment a therapeutically effective amount of a topoisomerase inhibitor topotecan, or a pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, and a therapeutically effective amount of thalidomide, or a pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof.
- 2. (Currently amended) A method of treating metastatic cancer which comprises administering to a patient in need of such treatment a therapeutically effective amount of a topoisomerase inhibitor topotecan, or a pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, and a therapeutically effective amount of thalidomide, or a pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof.
- 3. (Original) The method of claim 1 or 2 wherein the cancer is cancer of the head, neck, eye, mouth, throat, esophagus, chest, bone, lung, colon, rectum, stomach, prostate, breast, ovaries, kidney, liver, pancreas, and brain.
  - 4. (Original) The method of claim 3 wherein the cancer is colon or rectal cancer.
  - 5-7. (Canceled).
- 8. (Currently amended) The method of claim 7 1 or 2, wherein the irinotecan or SN 38, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 1 to about 1000 mg/m<sup>2</sup>, and the thalidomide, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 1 to about 2000 mg.

Application No. 09/853,617 Page 3 of 6

- 9. (Currently amended) The method of claim 8 wherein the irinotecan or SN-38, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 25 to about 750 mg/m<sup>2</sup>, and the thalidomide, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 50 to about 1000 mg.
- 10. (Currently amended) The method of claim 9 wherein the irinotecan or SN-38, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 50 to about 500 mg/m<sup>2</sup>, and the thalidomide, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 100 to about 750 mg.
- 11. (Currently amended) The method of claim 10 wherein the irinotecan or SN-38, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 100 to about 350 mg/m<sup>2</sup>, and the thalidomide, or pharmaceutically acceptable prodrug, salt, solvate, hydrate, or clathrate thereof, is administered in an amount of from about 200 to about 500 mg.

12-60. (Canceled).

- 61. (Previously presented) The method of claim 1 or 2, wherein thalidomide is administered.
- 62. (Previously presented) The method of claim 1 or 2, wherein the thalidomide salt or solvate is administered.